

Florida State University Libraries

2007-11-01

A review of web-assisted interventions for diabetes management: maximizing the potential for improving health outcomes.

Linda Lockett Brown, Mia Liza A Lustria and Jenice Rankins

This NIH-funded author manuscript originally appeared in PubMed Central at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2769687>.



A Review of Web-Assisted Interventions for Diabetes Management: Maximizing the Potential for Improving Health Outcomes

Linda Lockett Brown, M.Ag, RD, LD/N,¹ Mia Liza A. Lustria, Ph.D.,²
and Jenice Rankins, Ed.D.,¹

Abstract

Current endeavors in diabetes care focus on helping patients and providers deal successfully with the complexities of the disease by improving the system of care, expanding the reach of interventions, and empowering patients to engage in self-care behaviors. Internet technologies that combine the broad reach of mass media with the interactive capabilities of interpersonal media provide a wide range of advantages over standard modes of delivery. The technical affordances of Web delivery enable individualization or tailoring, appropriately timed reinforcement of educational messages, social support, improved feedback, and increased engagement. In turn, these have been significantly correlated with improved health outcomes.

This article is a narrative review of Web-based interventions for managing type 2 diabetes published from 2000 to 2007 that utilize Web sites, Web portals, electronic medical records, videoconference, interactive voice response, and short messaging systems. The most effective systems link medical management and self-management. Patient satisfaction is highest when the Web-based system gives them the ability to track blood glucose, receive electronic reminders, schedule physician visits, email their health care team, and interact with other diabetic patients. However, comprehensive medical and self-management programs have not been implemented widely outside of systems funded by government agencies. The cost of developing and maintaining comprehensive systems continues to be a challenge and is seldom measured in efficacy studies. Lack of reimbursement for Web-based treatments is also a major barrier to implementation. These barriers must be overcome for widespread adoption and realization of subsequent cost savings.

J Diabetes Sci Technol 2007;1(6):892-902

Author Affiliation: ¹College of Human Sciences, Florida State University, Tallahassee, Florida, and ²College of Information, Florida State University, Tallahassee, Florida

Abbreviations: (EMR) electronic medical records, (IVR) interactive voice response, (SMS) short messaging systems, (VA) Veterans Administration

Keywords: chronic care model, chronic disease, diabetes self-management, EMR, hemoglobin A1c, IVR, medical management, SMS, type 2 diabetes, videoconferencing, Web portal, Web site

Corresponding Author: Dr. Mia Liza A. Lustria, Assistant Professor, College of Information, 270 Louis Shores Building, Florida State University, Tallahassee, FL 32306-2100; email address mlustria@ci.fsu.edu

Introduction

The future of diabetes care lies in finding ways to help patients and providers deal successfully with the complexities of the disease by improving the system of care, expanding the reach of interventions, and empowering patients to engage in self-care behaviors.¹ Web-assisted diabetes management refers to interventions that involve the use of Internet technologies such as email, chat rooms, discussion groups, data uploads, electronic medical records, and short messaging systems (SMS) to enhance patient education, communication with providers, monitoring (e.g., blood glucose, diet, physical activity, blood pressure), goal setting, medication management, problem solving, and psychosocial adaptation. Web-assisted diabetes management can facilitate patient and physician roles to improve clinical outcomes.²

Internet technologies that marry the broad reach of mass media with the interactive capabilities of interpersonal media provide a wide range of advantages over standard modes. These include interactivity,^{3,4} homophily or common experience,⁵ social distance, and sense of privacy.⁶ These technical affordances enable individualization or tailoring,⁷⁻¹³ appropriately timed reinforcement of educational messages,^{14,15} social support,^{5,16-24} improved feedback between patient and provider,¹⁴ and increased engagement.²⁵ In turn, these have been significantly correlated with improved health outcomes.^{26,27} The United Kingdom Prospective Diabetes Study demonstrated that the risk for macrovascular and microvascular complications is reduced 14–37% with each percentage point decrease in hemoglobin A1c.²⁸

Diabetes management/care supplemented or delivered through Web technologies has been shown to decrease morbidity and mortality in a number of studies.^{4,29-32} Ultimately, these can lead to significant decreases in health care costs. Unfortunately, the tendency has been to design interventions around the technology rather than use the technology to meet core clinical or self-management outcomes. This article presents a narrative review of Web-based interventions for managing type 2 diabetes published from 2000 to 2007. Key issues affecting health outcomes, efficacy, design characteristics, application to diabetes management, and care are discussed. Finally, gaps in this area of research are described. The overall goal is to help shed light on how Web delivery contributes to improved health outcomes in diabetes care and how the use of Internet modalities can be maximized.

How Internet Technologies Have Been Used in Diabetes Treatment

A number of Internet and emerging Web technologies have been used to facilitate the delivery of diabetes care. These include Web sites, Web portals, electronic medical records (EMR), videoconference, interactive voice response (IVR), and SMS.

Web Site

Web sites are the most frequently used technology in diabetes care. Web sites are used to monitor, educate, provide feedback, and facilitate communication with medical staff. Patients have accessed Web sites via personal computers, messaging devices connected to plain old telephone system lines,^{33,34} cell phones,³⁵ and handheld devices.³⁶ A number of Web sites focus on diabetes self-management,^{33,34,37,38} weight management,^{36,39,40} physical activity,^{37,38,41,42} tailored physical activity,^{13,38,43} diet,^{13,38,44} blood glucose monitoring,^{13,35} data upload of blood glucose values,^{33,34,36,42} cardiovascular disease risk,^{36,45-47} and blood pressure.^{36,48}

Interactive Web site features increase knowledge, support engagement, increase self-efficacy, and facilitate behavior change through personalized feedback,^{38,43} resource libraries,^{42,45,46} tailored information,^{42,49} asynchronous discussion boards,⁴² synchronous chat,⁴² email to or from medical staff,^{38,42} and social support.^{18,38,44,50-52} Outcomes of trials lasting 12–120 weeks using Web sites showed a 1.2–16.0% decrease in hemoglobin A1c with 0–43% attrition reported. This is consistent with the meta-analysis reported by Norris and colleagues in 2002.¹⁵

The Veterans Administration (VA)^{34,36} and IDEATel Centers for Medicare and Medicaid Services demonstration projects are examples of comprehensive Web sites used for chronic disease management.^{45,46, 52-55} The VA program contained coordinated medical and self-management components. It is widely accepted that a combination of medical and self-management maximizes outcomes.⁵⁶ However, Meigs and co-workers⁵⁷ achieved a 2.7% drop in hemoglobin A1c with medical management alone. The intervention utilized a Web-based decision support tool developed to improve evidence-based management of type 2 diabetes. This is important because it highlights the critical role of getting the right information to physicians to promote implementation of evidence-based guidelines.⁵⁷⁻⁶¹ Web-based communication can facilitate this process.

Web sites have also been used successfully to screen for diabetic retinopathy⁶²⁻⁶⁴ and to monitor chronic wounds.^{65,66}

Web Portal

Web portals serve as points of access to information on the Internet from varied sources and are organized and presented in a cohesive manner.⁶⁷ Web portals have been used in diabetes treatment to (a) allow access to electronic medical records, medication, and health history; (b) provide preventive health care reminders, educational materials, and self-management resources, and (c) facilitate data uploads.^{45,46,68} In one study, patients rated the ability to log and track blood glucose levels over a 90-day period as one of the most useful features of a portal.⁶⁸ IDEATel used a Web portal for uploading and entry of clinical data.^{45,46} The 52-week intervention involving a large sample of 1665 patients in a self-management intervention achieved a 5.2% drop in hemoglobin A1c with an attrition rate of 18.2%.

There is some indication that portals have greater appeal for men, although women tended to access portals more often.⁶⁸ While portals are very popular, focus groups have shown consistently that patients are unwilling to pay for these services as they see the Internet as "free."⁶⁸⁻⁷¹

Electronic Medical Records

A number of studies have begun to look at how EMR's can be integrated with diabetes management. The patient's electronic medical record is the hub of the system. A feature common to most systems reviewed is the ability of patients and medical staff to enter data into the record.^{68,72,73} The goal of these systems is to improve disease outcomes by linking home-based monitoring, self-management support tools, case management, clinical information, and decision support systems.^{72,73} However, the technology must be integrated into a comprehensive disease management program for it to be effective. Use of the EMR alone to increase physicians' adherence to evidence-based practice guidelines is unlikely.^{59,74}

A word of caution is in order. It is imperative that a feedback loop is included in any system that allows direct interface between home-based monitoring systems and the EMR. Clinicians need to be alerted when values exceed certain thresholds so that a rapid response is enabled and treatment regimens can be altered.^{27,66,72-77}

Videoconferencing Systems

Videoconferencing involves the transmission of live video and audio between different locations. A growing

body of data suggests that videoconferencing used in diabetes care may be more effective than voice-only contact.^{45,46} The home telemedicine unit used in the IDEATel project included capabilities for synchronous videoconferencing with nursing staff at 8-15 frames per second. Results showed that the use of videoconferencing mimicked face-to-face encounters and increased patient satisfaction.⁶⁶

Interactive Voice Response Systems

Interactive voice response systems send patients recorded messages. Patients then respond by reporting clinical or other information using the telephone's touch-tone keypad, voice, or in-home messaging device.⁷⁸ This technology has been used successfully to decrease health care utilization costs for elderly diabetic patients in the VA system.³⁶ The intervention resulted in a 40% decrease in office and emergency room visits and a 30% drop in days hospitalized. Ironically, there was also a 16% increase in outpatient visits, which may have prevented more costly emergency room visits or in-patient hospitalization.³³

Short Messaging Systems

Short messaging systems send short text messages to or from mobile phones. Several interventions used SMS as part of an Internet-based blood glucose monitoring system. Patients uploaded blood glucose or blood pressure data to a secure Internet server via cell phone, which were immediately posted to their EMR. Medical staff reviewed and transmitted recommendations back to the patient via SMS.^{35,79}

One of the greatest challenges in diabetes management is maintenance of improved clinical outcomes. A 30-month study of a Web-based SMS showed that monitoring can reduce and maintain decreases in hemoglobin A1c. Levels were 7.7 ± 1.3 at baseline, 6.9 ± 1.1 at 15 months, and 6.7 ± 0.9 at 30 months.⁸⁰ The pervasiveness of cellular phones worldwide makes this type of system a very viable option. The caveat is having staff assigned to provide routine monitoring and feedback.^{72,81}

Characteristics of Web-Assisted Interventions

The following section discusses common characteristics of Web-assisted interventions. Web-assisted medical management refers to interventions that use technology to facilitate diagnosis, monitoring, and treatment. Web-assisted self management interventions focus on patient self-management or health education.

Medical Management

Adherence to evidence-based guidelines improves clinical outcomes for diabetes,⁵⁶ hypertension,⁸² and hyperlipidemia.⁸³ Systems that resulted in greater adherence to evidence-based guidelines included one or more of the following features: linking to electronic medical records,^{35,48,52,66,77,80} computerized prompts for staff and/or patients,^{84,85} reports, feedback, decision support,^{57,72,73} emailed clinical recommendations,⁵⁹ and electronic scheduling.^{57,66,85} Systems must be integrated seamlessly so that the staff perceive it as the way care is provided rather than an “extra” burden.⁵⁷

Self-Management

Web-assisted self-management interventions employed mechanisms for sustaining participation and provided personalized, immediate, or frequent feedback.^{35,38,53,72,80,86–91} The issue of attrition has been addressed in a number of ways, with significant improvements in retention when the intervention was provided as an adjunct to routine care⁹² or when interactivity increased.^{93,94} Stage-based interventions have been found to increase physical activity,⁹⁵ whereas others found action-based messages and ethnically appropriate messages beneficial in increasing engagement.³⁸

Which Intervention Features Are Most Useful?

Patients across studies rated several features favorably. These included the ability to log and track blood glucose values,^{13,28,35,66,96–98} receive electronic reminders,^{63,66,72,74,85,99} schedule physician visits,^{63,72,85} access relevant disease-specific information,^{65,74,99} email the health care team,^{66,68,85,96,98} get quick responses via email from staff,^{68,72,99} and form special interest groups with other members.^{38,68}

Key Issues

A number of recurring observations emerged from the studies reviewed. Ultimately, the goal of these interventions was to help patients reach clinical targets. The overarching challenge that seemed to emerge from this review was the limited ability to determine which intervention features are most effective.

Research Design

Sample sizes tended to be small and may not represent the populations in question adequately, which limits the generalizability of results. However, it is encouraging that ethnic minorities,^{38,39,46} Medicare recipients,^{33,46} and the elderly^{42,52,55} participated at a rate comparable to

traditional interventions.¹⁴ Few studies presented long-term results. Sixty percent of the studies were less than 6 months in length. It is important to determine if short-term results can be maintained over time.

Decreased Use

Web site use often decreased over time. It has been shown fairly consistently that improvement in clinical outcomes is directly proportional to dose.^{14,100} A closely related issue is attrition.

Attrition

Technical difficulties experienced by patients have been shown to be a major contributor to attrition.^{48,101} To address this, participants require assistance in overcoming technical issues early in the intervention and need to be monitored closely. Rewarding participants with gift certificates upon completion of tasks has successfully decreased attrition.³⁸

Self-Selection

Patients who choose to participate in programs tend to be more computer literate and more highly educated.^{68,71} This brings to the fore the question of whether people who are in most need of these interventions are being reached by these programs or whether Web delivery really improves access.¹⁰² However, the Diabetes-Net and IdeaTel projects overcame this barrier and successfully recruited participants that were representative of their target populations. Participants were recruited from the office of their primary care physician or were invited to participate by their primary care physician.^{44,46}

Behavior Change Framework

A recent meta-analysis of diabetes self-management interventions found that only one-third used a behavioral theory or model.¹⁰³ Programs based on sound behavioral theory have an increased probability of maximizing the effective application of Web-based technology.^{14,50,104–111}

Sustainability at the Organizational Level

Web-assisted interventions found in current diabetes management literature may have limited applicability to clinical settings where most care is provided.¹¹² Major projects tend to be government funded. Mechanisms to implement and sustain these systems in different practice settings need to be developed if widespread adoption is to occur.^{66–68,112}

The Chronic Care Model as described by Wagner and colleagues¹¹³ may provide a solution. The model contains six interdependent elements that serve to meet provider and patient needs with the goal of adherence to established clinical guidelines. The elements are evidence-based guidelines, practice redesign, patient education, expert system, and information. Each element can be facilitated by Web-based systems. Many features of the model overlap. Evidence-based guidelines have been used to generate clinical reminders for medical staff, design automated feedback, notify clinical staff values requiring timely intervention, and form the basis of treatment and/or decision support algorithms. Practice redesign most often shifted roles from physician to nurse and from clerical staff to an automated system. The net effect is that physicians can spend less time looking for data and have clinical guidelines readily available. Web-based patient education was the component of the model used most often. Numerous interventions required active patient participation for uploading data, responding to IVR, or entering information into diaries. The expert system element was the least-used component of the model. It is the most expensive to design and implement. The information element of the model utilized email, SMS, and telephone to contact patients.

Adoption of Chronic Care Model elements has been shown to improve the quality of care and increase the sustainability of comprehensive diabetes management programs.^{30,36,50,88,99,114–119} Implementation of a single model element can improve clinical outcomes in patients suffering from chronic illness.¹²⁰

Table 1 summarizes how elements of the model have been utilized in diabetes management interventions. The chronic care model was not used to structure the majority of the interventions listed. However, intentional application of model elements in the design of Web-based diabetes interventions may potentially increase adoption and improve the quality of care and clinical outcomes.^{117–120}

A case in point follows: a successful hospital-based diabetes management program operating since 1997 attributes three factors to its sustainability.⁸⁸ First, administrators took a long-term view of the program. Cost savings were not expected immediately. Second, processes were restructured to utilize staff and facilities effectively. Third, evidence-based guidelines were used to structure care delivery. The last two items are consistent with the chronic care model.^{113,117–119}

Barriers to Implementation

Barriers to implementation of Web-based medical or self-management interventions in diabetes occur at several levels. Five major types of barriers must be overcome. First, cost is frequently cited as a barrier to design, implementation, and sustainability of Web-based systems and needs to be examined more thoroughly.²⁷

The second barrier is failure to develop interventions that consider patient or staff characteristics in the design phase. These characteristics may have moderating effects on participation in Web-assisted care, clinical outcomes, or sustainability. Examples include functional literacy,¹²¹ health literacy,¹²² eHealth literacy,¹²³ computer literacy of the medical staff,¹²⁴ patient computer literacy,⁵³ cultural relevance,^{125–128} patient expectations,^{111,129} patient satisfaction,¹¹⁴ and provider satisfaction.¹³⁰

Third, evidence-based guidelines improve clinical outcomes. However, current medical practices and hospitals may require a redesign of internal processes to facilitate implementation of the guidelines. Examples from the studies reviewed are decision support systems or prompts delivered as orders are written^{48,57,85} or delivered when instructions are given to the patient.⁷⁷ Computer-based systems have the ability to seamlessly integrate clinical monitoring, medical records, medical management, and patient education.^{57,62–64,67,131,132}

Fourth, patients and medical professionals express concern about confidentiality. These concerns have limited the acceptance of Web-based communication, particularly email with patients.¹³³ Data privacy and security were cited in a recent review as barriers to implementation of Web-based chronic disease programs.²⁷

Fifth, payment policies must be changed to make disease management services billable. In most cases, physicians and facilities are unable to bill for time spent on Web-based diabetes care. Financial incentives to participate in these activities are lacking.^{134–136} Legislation is currently under consideration at the federal level that would remove this barrier from Medicare.¹³⁵ There is some indication that payment for these activities can be recouped in system-wide health care cost savings.¹³⁶

Areas for Future Study

Future studies need to answer a fundamental question: “Which technology is most effective for which person, under what circumstances, and why?”^{103,104} While we

Table 1.
Chronic Care Model Features Utilized in Interventions Delivered Via the Web

Elements ^a		Model examples ^a	Web interventions examples
1	Evidence-based guidelines		Standards of care for diabetes ⁵⁶ JNC 7 high blood pressure ⁸² National cholesterol guidelines ⁸³
2	Practice redesign	Appointments Roles Follow-up	High-risk patients selected from pharmacy database ⁵⁸ Computerized registry used to select high-risk patients ⁵⁹ Nursing staff case management ^{33–36,45,46,55,72,78,86,89,96,98,99} Retinopathy screening or wound monitoring ^{62–65} Staff prompted when EMR registers out-of-range hemoglobin A1c ⁷⁷ Clinical staff emails patients ¹³⁰ Patient phoned when uploaded data out of range ^{86,131} Reports generated for MD use during patient visit ^{48,101} MD sets clinical targets for patient alarms ⁴⁸
3	Patient education	Self-management Behavioral change Psychosocial support Patient participation	Patients transmit data into EMR ^{32,39,45,46,55,72,73,78,89,96,98} Patients access website information ^{35,38,42,43,44,45,46,55,89,131} Low literacy diabetes education ¹²¹ Tailored self care ^{13,40} Self-management support group ^{17,18,24,38,42,44,73} Electronic diary to record food intake and/or physical activity ^{42,72,131}
4	Expert system	Provider education Decision support Consultation	Clinical decision support tool ^{48,57} Staff receive and evaluate uploaded data ^{32,35,39,45,46,55,79,80,89,97,131} Clinical recommendations emailed to physician based on patient data ^{32,59} Computer-generated feedback to patient after data upload ⁷³ Automated SMS messages from data upload ⁹³ Care reminders generated from EMR data and delivered for physician use at point of care ^{84,85}
5	Information	Reminders Outcomes Feedback Care Planning	Physician emailed clinical recommendations based on evidence-based guidelines ⁵⁹ Videoconference for care planning and feedback ^{45,46,55,78,96} Web-based screening for retinopathy and wound monitoring ^{62–65} Point of care reminders ^{77,101,184} Patients receive SMS feedback about uploaded data ^{35,79,80,97,131} Patients receive emailed feedback about uploaded data ^{38,42,89} Patients emailed nursing staff ^{72,89}

^a Adapted from Wagner *et al.*¹¹³

have made progress in understanding important issues in designing Web-based diabetes management, there are still a number of areas that need further empirical investigation. Recommendations will be made in the areas of behavior change, effect of individual characteristics, selection of appropriate technology, and cost-effectiveness.

Theoretical Frameworks

The ultimate goal of diabetes care is behavior change that produces improved clinical outcomes. Structuring interventions based on sound behavioral theory will increase the likelihood of success.^{79,82,104} Frameworks for technology adoption also need to be applied to these systems.^{137,138}

Effect of Individual Characteristics

Further study is also needed to determine which individual characteristics affect use and level of engagement with Web-based interventions.^{24,50,51,54,103,104} These include psychosocial variables,^{52,90,108,109,140} level of adherence,^{106,107,131} literacy,^{41,121–123} age,^{46,52} ethnicity,^{38,39,58,71,102,110,125–128} determinants of satisfaction that impact attrition,^{14,104,129} self-efficacy,^{13,37,121} baseline hemoglobin A1c,^{42,34,36} comorbid conditions,^{32–34,42,139–141} and depression.^{44–47,55,105,140}

Selection of Appropriate Technology

Technology should be selected to facilitate the goals of the intervention. The mechanics of this process need

further study. The incremental effect of support groups, chat, and discussion boards on behavior change remains poorly understood.^{18,82,83,105,106} This was the focus of a well-designed study by Glasgow and colleagues⁴⁴ where the incremental effect of adding a personal coach and/or social support on clinical outcomes was explored. Clarification of how technological features affect outcomes in specific populations or enhance provider adoption will ultimately decrease development costs.¹³⁴⁻¹³⁹

Cost-Effectiveness

Methods to assess program development costs and overall cost-effectiveness need to be designed and validated. This will facilitate accurate comparison of studies. Health services research protocols may be useful in determining the cost-effectiveness of Web-based interventions.¹³⁶ Significant clinical outcomes and cost savings have been demonstrated in European systems that have successfully linked Web-based components to comprehensive disease management programs.¹⁰¹

Conclusion

Technological capability has increased rapidly. However, well-designed trials to determine how best to use the technology and that address public policy have not progressed at the same rate. Results from randomized controlled trials and carefully designed, nonrandomized trials can provide valuable information. Comprehensive systems that link medical management to self-management with the electronic medical record have consistently shown significant improvement in clinical outcomes and cost savings. These systems, however, have not been adopted widely.

Contrary to common beliefs about inertia with regards to patients' adoption of diabetes care technology, patients generally comply with expectations to collect and upload clinical data. This is consistent in older adults, across countries, socioeconomic levels, and ethnicities. However, clinicians have not always responded to uploaded EMR data with timely review and feedback. In this regard, programs have been found to work best when staff is specifically assigned to support Web-assisted interventions. Design, implementation, and maintenance of effective Web-based diabetes management systems may best be accomplished by the collaboration of patients, clinicians, engineers, private/public payers, and professionals skilled in facilitating organizational change. To foster this, the Chronic Care Model provides a workable framework for developing sustainable systems that meet the needs of both patients and medical staff.

References:

1. Vinicor F. The future of diabetes: what is there besides new medicines? *Clin Diabetes*. 2004;22(2):94-6.
2. Vincent D, Clark L, Zimmer LM, Sanchez J. Using focus groups to develop a culturally competent diabetes self-management program for Mexican Americans. *Diabetes Educ*. 2006;32(1):89-97.
3. Glasgow RE, Bull S. Making a difference with interactive technology: considerations in using and evaluating computerized aids for diabetes self-management education. *Diabetes Spectrum*. 2001;14(2):99-106.
4. Jackson CL, Bolen S, Brancati FL, Batts-Turner ML, Gary TL. A systematic review of interactive computer-assisted technology in diabetes care: interactive information technology in diabetes care. *J Gen Intern Med*. 2006;21(2):105-10.
5. McKay HG, Feil EG, Glasgow RE, Brown JE. Feasibility and use of an internet support service for diabetes self-management. *Diabetes Educ*. 1998;24(2):174-9.
6. Walther JB, Pingree S, Hawkins RP, Buller DB. Attributes of interactive online health information systems. *J Med Internet Res*. 2005;7(3):e33.
7. Brug J, Oenema A, Campbell M. Past, present, and future of computer-tailored nutrition education. *Am J Clin Nutr*. 2003;77(4):1028S-1034S.
8. Glasgow RE. Translating research to practice: lessons learned, areas for improvement, and future directions. *Diabetes Care*. 2003;26(8):2451-6.
9. Glasgow RE, Toobert DJ. Brief, computer-assisted diabetes dietary self-management counseling: effects on behavior, physiologic outcomes, and quality of life. *Med Care*. 2000;38(11):1062-73.
10. Glazier RH, Bajcar J, Kennie NR, Willson K. A systematic review of interventions to improve diabetes care in socially disadvantaged populations. *Diabetes Care*. 2006;29(7):1675-88.
11. Oenema A, Brug J. Feedback strategies to raise awareness of personal dietary intake: results of a randomized controlled trial. *Prev Med*. 2003;36(4):429-39.
12. Suggs LS. A 10-year retrospective of research in new technologies for health communication. *J Health Comm*. 2006;11(1):61-74.
13. Wangberg SC. An Internet-based diabetes self-care intervention tailored to self-efficacy. *Health Educ Res*. 2007 Apr 5; [Epub ahead of print]
14. Norris SL, Engelgau MM, Venkat Narayan KM. Effectiveness of self-management training in type 2 diabetes: a systematic review of randomized controlled trials. *Diabetes Care*. 2001;24(3):561-87.
15. Norris SL, Lau J, Smith SJ, Schmid CH, Engelgau MM. Self-management education for adults with type 2 diabetes: a meta-analysis of the effect on glycemic control. *Diabetes Care*. 2002;25(7):1159-71.
16. Loader BD, Muncer S, Burrows R, Pleace N, Nettleton S. Medicine on the line? Computer-mediated social support and advice for people with diabetes. *Int J Soc Welfare*. 2002;11(1):53-65.
17. Barrera M, Glasgow RE, McKay HG, Boles SM, Feil EG. Do Internet-based support interventions change perceptions of social support? An experimental trial of approaches for supporting diabetes self-management. *Am J Comm Psychol*. 2002;30(5):637-54.
18. Barrera M Jr, Toobert DJ, Angell KL, Glasgow RE, MacKinnon DP. Social support and social-ecological resources as mediators of lifestyle intervention effects for type 2 diabetes. *J Health Psychol*. 2006;11(3):483-95.
19. DiMatteo MR. Social support and patient adherence to medical treatment: a meta-analysis. *Health Psychol*. 2004;23(2):207-18.

20. Peyrot M, Rubin RR, Siminerio LM. Physician and nurse use of psychosocial strategies in diabetes care: results of the cross-national diabetes attitudes, wishes and needs (DAWN) study. *Diabetes Care*. 2006;29(6):1256-62.
21. Powell JA, Lowe P, Griffiths FE, Thorogood M. A critical analysis of the literature on the Internet and consumer health information. *J Telemed Telecare*. 2005;11(Suppl.1):41-3.
22. White M, Dorman SM. Receiving social support online: implications for health education. *Health Educ Res*. 2001;16(6):693-707.
23. Zhang X, Norris SL, Gregg EW, Beckles G. Social support and mortality among older persons with diabetes. *Diabetes Educ*. 2007;33(2):273-81.
24. Zrebiec JF, Jacobson AM. What attracts patients with diabetes to an Internet support group? A 21-month longitudinal website study. *Diabet Med*. 2001;18(2):154-8.
25. Gomaa WH, Morrow T, Muntendam P. Technology-based disease management: a low-cost, high-value solution for the management of chronic disease. *Dis Manag Health Outcomes*. 2001;9(10):577-88.
26. Norris SL, Nichols PJ, Caspersen CJ, Glasgow RE, Engelgau MM, Jack L, Isham G, Snyder SR, Carande-Kulis VG, Garfield S, Briss P, McCulloch D. The effectiveness of disease and case management for people with diabetes. A systematic review. *Am J Prev Med*. 2002;22(4):15-38.
27. Dorr D, Bonner LM, Cohen AN, Shoai RS, Perrin R, Chaney E, Young AS. Informatics systems to promote improved care for chronic illness: a literature review. *J Am Med Inform Assoc*. 2007;14(2):156-63.
28. Stratton IM, Adler AI, Neil HA, Matthews DR, Manley SE, Cull CA, Hadden D, Turner RC, Holman RR. Association of glycaemia with macrovascular and microvascular complications of type 2 diabetes (UKPDS 35): prospective observational study. *BMJ*. 2000;321(7258):405-12.
29. Gimenez-Perez G, Caixas A, Gimenez-Palop O, Gonzalez-Clemente JM, Mauricio D. Dissemination of 'patient-oriented evidence that matters' on the Internet: the case of type 2 diabetes treatment. *Diabet Med*. 2005;22(6):688-92.
30. Glasgow RE, Hiss RG, Anderson RM, Friedman NM, Hayward RA, Marrero DG, Taylor CB, Vinicor F. Report of the health care delivery work group: behavioral research related to the establishment of a chronic disease model for diabetes care. *Diabetes Care*. 2001;24(1):124-30.
31. Mazzi C, Kiss M. A framework for the evaluation of Internet-based diabetes management. *J Med Internet Res*. 2002;4(1):e1.
32. Bellazzi R, Larizza C, Montani S, Riva A, Stefanelli M, d'Annunzio G, Lorini R, Gomez EJ, Hernando E, Bragues E, Cermeno J, Corcoy R, de Leiva A, Cobelli C, Nucci G, Del Prato S, Maran A, Kilkki E, Tuominen J. A telemedicine support for diabetes management: the T-IDDM project. *Comput Methods Programs Biomed*. 2002;69(2):147-61.
33. Dang S, Ma F, Nedd N, Aguilar E, Roos B. Differential resource utilization benefits with Internet-based care coordination in elderly veterans with chronic diseases associated with high resource utilization. *Telemed J E Health*. 2006;12(1):14-23.
34. Joseph AM. Care coordination and telehealth technology in promoting self-management among chronically ill patients. *Telemed J E Health*. 2006;12(2):156-9.
35. Kwon HS, Cho JH, Kim HS, Song BR, Ko SH, Lee JM, Kim SR, Chang SA, Kim HS, Cha BY, Lee KW, Son HY, Lee JH, Lee WC, Yoon KH. Establishment of blood glucose monitoring system using the Internet. *Diabetes Care*. 2004;27(2):478-83.
36. Chumbler N, Neugaard B, Kobb R, Ryan P, Qin H, Joo Y. An observational study of veterans with diabetes receiving weekly or daily home telehealth monitoring. *J Telemed Telecare*. 2005;11:150-6.
37. Lorig KR, Ritter PL, Laurent DD, Plant K. Internet-based chronic disease self-management. *Med Care*. 2006;44(11):964-71.
38. Robertson C, Kattelman KP, Ren CP. Control of type 2 diabetes mellitus using interactive Internet-based support on a Northern Plains Indian reservation: a pilot study. *Top Clin Nutr*. 2007;22(2):185-93.
39. Bondmass M, Bolger N, Castro G, Avitall B. The effect of home monitoring and telemanagement on blood pressure control among African Americans. *Telemed J*. 2000;6(1):15-23.
40. Kalten MR, Ardito DA, Cimino C, Wylie-Rosett J. A web-accessible core weight management program. *Diabetes Educ*. 2000;26(6):929-36.
41. Kim S, Love F, Quistberg DA, Shea JA. Association of health literacy with self-management behavior in patients with diabetes. *Diabetes Care*. 2004;27(12):2980-2.
42. Bond GE, Burr R, Wolf FM, Price M, McCurry SM, Teri L. Preliminary findings of the effects of comorbidities on a web-based intervention on self-reported blood sugar readings among adults age 60 and older with diabetes. *Telemed J E Health*. 2006;12(6):707-10.
43. McKay HG, King D, Eakin EG, Seeley JR, Glasgow RE. The diabetes network internet-based physical activity intervention: a randomized pilot study. *Diabetes Care*. 2001;24(8):1328-34.
44. Glasgow RE, Boles SM, McKay HG, Feil EG, Barrera M Jr. The D-Net diabetes self-management program: long-term implementation, outcomes, and generalization results. *Prev Med*. 2003;36:410-9.
45. Shea S, Starren J, Weinstock RS, Knudson PE, Teresi J, Holmes D, Palmas W, Field L, Goland R, Tuck C, Hripacsak G, Capps L, Liss D. Columbia University's Informatics for Diabetes Education and Telemedicine (IDEATel) project: rationale and design. *J Am Med Inform Assoc*. 2002;9(1):49-62.
46. Shea S, Weinstock RS, Starren J, Teresi J, Palmas W, Field L, Morin P, Goland R, Izquierdo RE, Wolff LT, Ashraf M, Hillman C, Silver S, Meyer S, Holmes D, Petkova E, Capps L, Lantigua RA. A randomized trial comparing telemedicine case management with usual care in older, ethnically diverse, medically underserved patients with diabetes mellitus. *J Am Med Inform Assoc*. 2006;13(1):40-51.
47. Kendall DM, Bergenstal RM. Comprehensive management of patients with type 2 diabetes: establishing priorities of care. *Am J Manag Care*. 2001;7(10):S327-43.
48. Luzio S, Piehlmeier W, Tovar C, Eberl S, Lätzsch G, Fallböhmer E, Rumpel E, Owens DR, Landgraf R. Results of the pilot study of DIADEM: a comprehensive disease management programme for type 2 diabetes. *Diabetes Res Clin Prac*. 2007;76:410-7.
49. Liu CT, Yeh YT, Lee TI, Li YC. Observations on online services for diabetes management. *Diabetes Care*. 2005;28(11):2807-8.
50. Glasgow R, Whitesides H, Nelson C, King D. Use of the patient assessment of chronic illness care (PACIC) with diabetic patients: relationship to patient characteristics, receipt of care, and self-management. *Diabetes Care*. 2003;26:55-61.
51. Glasgow RE, Toobert DJ, Barrera M Jr, Strycker LA. The Chronic Illness Resources Survey: cross-validation and sensitivity to intervention. *Health Educ Res*. 2005;20(4):402-9.

52. Palmas W, Teresi J, Morin P, Wolff LT, Field L, Eimicke JP, Capps L, Prigollini A, Orbe I, Weinstock RS, Shea S. Recruitment and enrollment of rural and urban medically underserved elderly into a randomized trial of telemedicine care management for diabetes care. *Telemed J E Health*. 2006;12(5):601-7.
53. Bakken S, Grullon-Figueroa L, Izquierdo R, Lee NJ, Morin P, Palmas W, Teresi J, Weinstock RS, Shea S, Starren J; IDEATel Consortium. Development, validation, and use of English and Spanish versions of the telemedicine satisfaction and usefulness questionnaire. *J Am Med Inform Assoc*. 2006;13(6):660-7.
54. Kaufman DR, Starren J, Patel VL, Morin PC, Hilliman C, Pevzner J, Weinstock RS, Goland R, Shea S. A cognitive framework for understanding barriers to the productive use of a diabetes home telemedicine system. *AMIA Annu Symp Proc*. 2003;356-60.
55. Trief PM, Morin PC, Izquierdo R, Teresi J, Eimicke JP, Goland R, Starren J, Shea S, Weinstock RS. Depression and glycemic control in elderly ethnically diverse patients with diabetes: the IDEATel Project. *Diabetes Care*. 2006;29(4):830-5.
56. American Diabetes Association. Standards of medical care in diabetes. *Diabetes Care*. 2007;30:54-541.
57. Meigs JB, Cagliero E, Dubey A, Murphy-Sheehy P, Gildesgame C, Chueh H, Barry MJ, Singer DE, Nathan DM. A controlled trial of web-based diabetes disease management: the MGH diabetes primary care improvement project. *Diabetes Care*. 2003;26(3):750-7.
58. Davidson MB. Effect of nurse-directed diabetes care in a minority population. *Diabetes Care*. 2003;26(8):2281-7.
59. Grant RW, Hamrick HE, Sullivan CM, Dubey AK, Chueh HC, Cagliero E, Meigs JB. Impact of population management with direct physician feedback on care of patients with type 2 diabetes. *Diabetes Care*. 2003;26(8):2275-80.
60. Herman WH. Evidence-based diabetes care. *Clin Diabetes*. 2002;20(1):22-3.
61. Piette JD. The future of diabetes disease management: integrating lessons learned from clinical, health services and policy research. *Am J Manag Care*. 2005;11(4):203-5.
62. Abramoff MD, Suttrop-Schulten MSA. Web-based screening for diabetic retinopathy in a primary care population: the EyeCheck project. *Telemed J E Health*. 2005;11(6):668-74.
63. Kawasaki S, Ito S, Satoh S, Mori Y, Saito T, Fukushima H, Kato S, Sekihara H. Use of telemedicine in periodic screening of diabetic retinopathy. *Telemed J E Health*. 2003;9(3):235-9.
64. Wei JC, Valentino DJ, Bell DS, Baker RS. A web-based telemedicine system for diabetic retinopathy screening using digital fundus photography. *Telemed J E Health*. 2006;12(1):50-7.
65. Kim MH, Lowery JC, Hamill J, Wilkins EG. Accuracy of a web-based system for monitoring chronic wounds. *Telemed J E Health*. 2003;9(2):129-39.
66. Noel HC, Vogel DC, Erdos JJ, Cornwall D, Levin F. Home telehealth reduces healthcare costs. *Telemed J E Health*. 2004;10(2):170-83.
67. Chau M, Huang Z, Qin J, Zhou Y, Chen H. Building a scientific knowledge web portal: The NanoPort experience. *Decision Support Systems*. 2006;42(2):1216-38.
68. Hess R, Bryce CL, McTigue K, Fitzgerald K, Zickmund S, Olshansky E, Fischer G. The diabetes patient portal: patient perspectives on structure and delivery. *Diabetes Spectr*. 2006;19(2):106-10.
69. Johnson F, Manjunath R, Mansfield C, CLayton L, Hoerger T, Zhang P. High-risk individuals' willingness to pay for diabetes risk-reduction programs. *Diabetes Care*. 2006;28(6):1351-6.
70. Dou W. Will Internet users pay for online content? *J Advertis Res*. 2004;44(4):349-59.
71. Jackson CL, Batts-Turner M, Falb MD, Yeh HC, Brancati FL, Gary TL. Computer and internet use among urban African Americans with type 2 diabetes. *J Urban Health*. 2005;82(4):575-83.
72. Goldberg HI, Ralston JD, Hirsch IB, Hoath JI, Ahmed KI. Using an Internet comanagement module to improve the quality of chronic disease care. *Jt Comm J Qual Saf*. 2003;29(9):443-51.
73. De Clercq P, Hasman A, Wolffenbuttel B. A consumer health record for supporting the patient-centered management of chronic diseases. *Med Inform Internet Med*. 2003;28(2):117-27.
74. Crosson JC, Ohman-Strickland PA, Hahn KA, DiCicco-Bloom B, Shaw E, Orzano AJ, Crabtree BF. Electronic medical records and diabetes quality of care: results from a sample of family medicine practices. *Ann Fam Med*. 2007;5(3):209-15.
75. Bellazzi R. Electronic management systems in diabetes mellitus: impact on patient outcomes. *Dis Manag Health Outcomes*. 2003;11(3):159-71.
76. Klonoff DC. Diabetes and telemedicine: is the technology sound, effective, cost-effective, and practical? *Diabetes Care*. 2003;26(5):1626-8.
77. Chima CS, Farmer-Dziak N, Cardwell P, Snow S. Use of technology to track program outcomes in a diabetes self-management program. *J Am Diet Assoc*. 2005;105:1933-8.
78. O'Connell MA, Cherry JC. The Health Hero Online Service: A new Internet-based communications platform for disease management and performance management. *Pract Dis Manag*. 2000;7(3):149-61.
79. Kim H, Jeong H. A nurse short message service by cellular phone in type-2 diabetic patients for six months. *J Clin Nurs*. 2007;16(6):1082-7.
80. Cho JH, Chang SA, Kwon HS, Choi YH, Ko SH, Moon SD, Yoo SJ, Song KH, Son HS, Kim HS, Lee WC, Cha BY, Son HY, Yoon KH. Long-term effect of the Internet-based glucose monitoring system on HbA1c reduction and glucose stability: a 30-month follow-up study for diabetes management with a ubiquitous medical care system. *Diabetes Care*. 2006;29(12):2625-31.
81. Hersh WR, Hickam DH, Severance SM, Dana TL, Krages KP, Helfand M. Diagnosis, access and outcomes: update of a systematic review of telemedicine services. *J Telemed Telecare*. 2006;12(Suppl 2):S23-31.
82. Chobanian AV, Bakvis GL, Black HR, Cushman WC, Green LA, Izzo JL Jr, Jones DW, Materson BJ, Oparil S, Wright JT Jr, Roccella EJ; National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA*. 2003;289:2560-72.
83. Grundy SM. Approach to lipoprotein management in 2001 National Cholesterol Guidelines. *Am J Cardiol*. 2002;90(8):11i-21i.
84. Sciamanna CN, Gifford DR, Smith RJ. Design and acceptability of patient-oriented computerized diabetes care reminders for use at the point of care. *Med Inform Internet Med*. 2004;29(2):157-68.
85. Sequist TD, Gandhi TK, Karson AS, Fiskio JM, Bugbee D, Sperling M, Cook EF, Orav EJ, Fairchild DG, Bates DW. A randomized trial of electronic clinical reminders to improve quality of care for diabetes and coronary artery disease. *J Am Med Inform Assoc*. 2005;12(4):431-7.
86. Piette JD, Weinberger M, Kraemer FB, McPhee SJ. Impact of automated calls with nurse follow-up on diabetes treatment outcomes in a department of Veterans Affairs health care system: a randomized controlled trial. *Diabetes Care*. 2001;24(2):202-8.

87. Chevront MM, Cassell C, Jackson B. Health communication on the internet: an effective channel for health behavior change? *J Health Comm.* 1998;3(1):71-9.
88. Whittaker SL, Adkins S, Phillips R, Jones J, Horsley MA, Kelley G. Success factors in the long-term sustainability of a telediabetes programme. *J Telemed Telecare.* 2004;10:84-8.
89. McMahon GT, Gomes HE, Hickson Hohne S, Hu TM, Levine BA, Conlin PR. Web-based care management in patients with poorly controlled diabetes. *Diabetes Care.* 2005;28(7):1624-9.
90. Strecher V. Internet methods for delivering behavioral and health-related interventions (eHealth). *Annu Rev Clin Psychol.* 2007;3:325-48.
91. Campbell MK, DeVellis BM, Strecher VJ, Ammerman AS, DeVillis RF, Sanchez RS. Improving dietary behavior: The effectiveness of tailored messages in primary care settings. *Am J Public Health.* 1994;84:783-7.
92. Kypri K, McAnally HM. Randomized controlled trial of a web-based primary care intervention for multiple health risk behaviors. *Prev Med.* 2005;41(3-4):761-6.
93. Hurling R, Catt M, Boni MD, Fairley BW, Hurst T, Murray P, Richardson A, Sodhi JS. Using internet mobile phone technology to deliver an automated physical activity program: randomized controlled trial. *J Med Internet Res.* 2007;9(2):e7.
94. Hurling R, Fairley BW, Dias MB. Internet-based exercise interventions: are more interactive designs better? *Psychol Health.* 2006;21(6):757-2.
95. Marshall AL, Leslie ER, Bauman AE, Marcus BH, Owen N. Print versus website physical activity programs: a randomized trial. *Am J Prev Med.* 2003;25(2):88-94.
96. O'Connell MA, Cherry JC. The Health Hero online service: A new Internet-based communications platform for disease management, case management and performance measurement. *Dis Manag Health Outcomes.* 2000;7(3):149-61.
97. Hee-Sung K. Impact of web-based nurse's education on glycosylated haemoglobin in type 2 diabetic patients. *J Clin Nurs.* 2007;16(7):1361-6.
98. Cherry JC, Moffatt T, Rodriguez C, Dryden K. Diabetes disease management program for an indigent population empowered by telemedicine technology. *Diabetes Technol Ther.* 2002;4(6):783-91.
99. Barnett TE, Chumblor NR, Vogel WB, Beyth RJ, Qin H, Kobb R. The effectiveness of a care coordination home telehealth program for veterans with diabetes mellitus: a 2-year follow-up. *Am J Manag Care.* 2006;2(8):467-74.
100. Norris SL, Zhang X, Avenell A, Gregg E, Bowman B, Serdula M, Brown TJ, Schmid CH, Lau J. Long-term effectiveness of lifestyle and behavioral weight loss interventions in adults with type 2 diabetes: a meta-analysis. *Am J Med.* 2004;117(10):762-74.
101. Salzseider E, Augstein P, Vogt L, Klaus-Dieter Kohnert, Heinke P, Freyse, EJ, Ahmed AA, Metwali Z, Salman I, Attef O. Telemedicine-based KADIS combined with CGMS has high potential for improving outpatient diabetes care. *J Diabetes Sci Technol.* 2007;1(4):511-21.
102. Chang BL. Internet intervention for community elders: process and feasibility. *West J Nurs Res.* 2004;26(4):461-6.
103. Welch G, Shayne R. Interactive behavioral technologies and diabetes self-management support: recent research findings from clinical trials. *Curr Diabetes Rep.* 2006;6:130-6.
104. Evers KE, Cummins CO, Prochaska JO, Prochaska JM. Online health behavior and disease management programs: are we ready for them? Are they ready for us? *J Med Internet Res.* 2005;7(3):e27.
105. Glasgow R, Toobert D, Barrera M Jr, Strycker LA. The chronic illness resources survey: cross-validation and sensitivity to intervention. *Health Educ Res.* 2005;20(4):402-9.
106. Schechter CB, Walker EA. Improving adherence to diabetes self-management recommendations. *Diabetes Spectrum.* 2002;15(3):170-5.
107. Walker EA. Health behavior: from paradox to paradigm. *Diabetes Spectrum.* 2001;14(1):6-8.
108. Vallis M, Ruggiero L, Greene G, Jones H, Zinman B, Rossi S, Edwards L, Rossi JS, Prochaska JO. Stages of change for healthy eating in diabetes: relation to demographic, eating-related, health care utilization, and psychosocial factors. *Diabetes Care.* 2003;26(5):1468-74.
109. Delamater AM, Jacobson AM, Anderson B, Cox D, Fisher L, Lustman P, Rubin R, Wysocki T; Psychosocial Therapies Working Group. Psychosocial therapies in diabetes: report of the Psychosocial Therapies Working Group. *Diabetes Care.* 2001;24(7):1286-92.
110. Mau MK, Glanz K, Severino R, Grove JS, Johnson B, Curb JD. Mediators of lifestyle behavior change in native Hawaiians: initial findings from the native Hawaiian diabetes intervention program. *Diabetes Care.* 2001;24(10):1770-5.
111. Wilson EV, Lankton NK. Modeling patients' acceptance of provider-delivered e-health. *J Am Med Inform Assoc.* 2004;11(4):241-8.
112. Leeman J. Interventions to improve diabetes self-management: utility and relevance for practice. *Diabetes Educ.* 2006;32(4):571-83.
113. Wagner EH, Austin BT, Von Korff M. Organizing care for patients with chronic illness. *Milbank Q.* 1996;4:511-44.
114. Glasgow RE, Wagner EH, Schaefer J, Mahoney LD, Reid RJ, Greene SM. Development and validation of the patient assessment of chronic illness care (PACIC). *Med Care.* 2005;43(5):436-44.
115. Siminerio LM, Piatt G, Zgibor JC. Implementing the chronic care model for improvements in diabetes care and education in a rural primary care practice. *Diabetes Educ.* 2005;31(2):225-34.
116. Chin MH, Cook S, Drum ML, Jin L, Guillen M, Humikowski CA, Koppert J, Harrison JF, Lippold S, Schaefer CT; Midwest cluster health disparities collaborative. Improving diabetes care in midwest community health centers with the health disparities collaborative. *Diabetes Care.* 2004;27(1):2-8.
117. Nutting PA, Dickinson WP, Dickinson LM, Nelson CC, King DK, Crabtree BF, Glasgow RE. Use of chronic care model elements is associated with higher-quality care for diabetes. *Ann Fam Med.* 2007;5(1):14-20.
118. Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness: the chronic care model, Part 2. *JAMA.* 2002;288:1909-14.
119. Bodenheimer T, Wagner EH, Grumbach K. Improving primary care for patients with chronic illness. *JAMA.* 2002;288:1775-9.
120. Tsai AC, Morton SC, Mangione CM, Keeler EB. A meta-analysis of interventions to improve care for chronic illness. *Am J Manag Care.* 2005;11:478-88.
121. Gerber BS, Brodsky IG, Lawless KA, Smolin LI, Arozullah AM, Smith EV, Berbaum ML, Heckerling PS, Eiser AR. Implementation and evaluation of a low-literacy diabetes education computer multimedia application. *Diabetes Care.* 2005;28(7):1574-80.
122. Zarcadoolas C, Pleasant A, Greer DS. Understanding health literacy: an expanded model. *Health Promot. Int.* 2005;20(2):195-203.
123. Norman CD, Skinner HA. eHealth literacy: essential skills for consumer health in a networked world. *J Med Internet Res.* 2006;8(2):e9.

124. Lyons SS, Tripp-Reimer T, Sorofman BA, Dewitt JE, Bootsmiller BJ, Vaughn TE, Doebbeling BN. VA QUERI Informatics paper: information technology for clinical guideline implementation: perceptions of multidisciplinary stakeholders. *J Am Med Inform Assoc.* 2005;12(1):64-71.
125. Anderson RM, Funnell MM, Arnold MS, Barr PA, Edwards GJ, Fitzgerald JT. Assessing the cultural relevance of an education program for urban African Americans with diabetes. *Diabetes Educ.* 2000;26(2):280-9.
126. Wu JH, Haan MN, Liang J, Ghosh D, Gonzalez HM, Herman WH. Diabetes as a predictor of change in functional status among older Mexican Americans: a population-based cohort study. *Diabetes Care.* 2003;26(2):314-9.
127. Alferi SM, Carver CS, Antoni MH, Weiss S, Duran RE. An exploratory study of social support, distress, and life disruption among low-income Hispanic women under treatment for early stage breast cancer. *Health Psychol.* 2001;20(1):41-6.
128. Lorig KR, Ritter PL, Gonzalez VM. Hispanic chronic disease self-management: a randomized community-based outcome trial. *Nurs Res.* 2003;52(6):361-9.
129. Stevenson FA, Cox K, Britten N, Dundar Y. A systematic review of the research on communication between patients and health care professionals about medicines: the consequences for concordance. *Health Expect.* 2004;7(3):235-45.
130. Katz SJ, Moyer CA, Cox DT, Stern DT. Effect of a triage-based E-mail system on clinic resource use and patient and physician satisfaction in primary care: a randomized controlled trial. *J Gen Intern Med.* 2003;18(9):736-44.
131. Kim HS, Oh JA. Adherence to diabetes control recommendations: impact of nurse telephone calls. *J Adv Nurs.* 2003; 44(3):231-4.
132. Chau D, Edelman SV. Clinical management of diabetes in the elderly. *Clin Diabetes.* 2001;19(4):172-5.
133. Bell DS, Daly DM, Robinson P. Is there a digital divide among physicians? A geographic analysis of information technology in southern California physician offices. *J Am Med Inform Assoc.* 2003;10(5):484-93.
134. Welch WP, Bazarko D, Ritten K, Burgess Y, Harmon R, Sandy LG. Electronic health records in four community physician practices: impact on quality and cost of care. *J Am Med Inform Assoc.* 2007;14(3):320-8.
135. Ubl SJ. Public policy implications for using remote monitoring technology to treat diabetes. *J Diabetes Sci Technol.* 2007;1(3):436-9.
136. Curtin K, Beckman H, Pankow G, Milillo Y, Green RA. Return on investment in pay for performance: a diabetes case study. *J Healthc Manag.* 2006;51(6):365-74.
137. Whitten P, Sypher BD. Evolution of telemedicine from an applied communication perspective in the United States. *Telemed J E Health.* 2006;12(5):590-600.
138. Glasgow RE, Nelson CC, Strycker LA, King DK. Using RE-AIM metrics to evaluate diabetes self-management support interventions. *Am J Prev Med.* 2006;30(1):67-73.
139. Glasgow RE. eHealth evaluation and dissemination research. *Am J Prev Med.* 2007;32(5) Suppl 1:S119-26.
140. Taylor CB, Luce KH. Computer- and Internet-based psychotherapy interventions. *Curr Direct Psychol Sci.* 2003;12(1):18-22.
141. Zhou H, Isaman DJ, Messinger S, Brown MB, Klein R, Brandle M, Herman WH. A computer simulation model of diabetes progression, quality of life, and cost. *Diabetes Care.* 2005;28(12):2856-63.